

The one-sided kissing number in four dimensions

Oleg R. Musin *

Abstract

Let H be a closed half-space of n -dimensional Euclidean space. Suppose S is a unit sphere in H that touches the supporting hyperplane of H . The one-sided kissing number $B(n)$ is the maximal number of unit nonoverlapping spheres in H that can touch S . Clearly, $B(2) = 4$. It was proved that $B(3) = 9$. Recently, K. Bezdek proved that $B(4) = 18$ or 19, and conjectured that $B(4) = 18$. We present a proof of this conjecture.

1 Introduction

Let H be a closed half-space of $\mathbf{R}^n$. Suppose S is a unit sphere in H that touches the supporting hyperplane of H . The one-sided kissing number $B(n)$ is the maximal number of unit nonoverlapping spheres in H that can touch S .

The problem of finding $B(3)$ was raised by L. Fejes Tóth and H. Sachs in 1976 [11]. K. Bezdek and P. Brass [4] studied the problem in more general setting and they introduced the term “one-sided Hadwiger number”, which in the case of a ball is the same as the one-sided kissing number. The term “one-sided kissing number” have been introduced by K. Bezdek [2, 3].




Fig. 1

The one-sided kissing problem looks more complicated than the classical kissing number problem. Actually, that is a constrained kissing problem. Denote by $k(n)$ the kissing number in n -dimensions, i.e. $k(n)$ is the highest number of equal nonoverlapping spheres in $\mathbf{R}^n$ that can touch another sphere of the same size. Currently, $k(n)$ known for $n = 2, 3, 4, 8, 24$: $k(2) = 6$, $k(3) = 12$

*Institute for Math. Study of Complex Systems, Moscow State University, Moscow, Russia
omusin@mail.ru

[22], $k(4) = 24$ [15, 17], $k(8) = 240$, $k(24) = 196, 560$ [14, 18]. There are strong relations between $k(n)$ and $B(n)$ (see the Conclusions and Theorem 1 in this paper).

If M unit spheres kiss (touch) the unit sphere in $H \subset \mathbf{R}^n$, then the set of kissing points is an arrangement on the hemisphere S_+ of S such that the (Euclidean) distance between any two points is at least 1. So the one-sided kissing number problem can be stated in another way: How many points can be placed on the surface of S_+ so that the angular separation between any two points is at least $\pi/3$?

Clearly, $B(2) = 4$ (see Fig. 1). It have been proved by G. Fejes Tóth [9] in 1981 that $B(3) = 9$ (see also H. Sachs [20] and A. Bezdek & K. Bezdek [1] for other proofs). G. Kertész (1994) proved that the maximal one-sided kissing arrangement is unique (up to isometry) on the 2-hemisphere S_+ .

Note that $B(4) \geq 18$. Indeed, let $A = 1/\sqrt{2}$,

$$\begin{aligned} p_1 &= (A, 0, 0, A), & p_2 &= (-A, 0, 0, A), \\ p_{\{3,4\}} &= (0, \pm A, 0, A), & p_{\{5,6\}} &= (0, 0, \pm A, A), \\ p_{\{7,\dots,10\}} &= (\pm A, \pm A, 0, 0), & p_{\{11,\dots,14\}} &= (\pm A, 0, \pm A, 0), \\ p_{\{15,\dots,18\}} &= (0, \pm A, \pm A, 0). \end{aligned}$$

Then these 18 points $\{p_i\}$ lie on the closed upper hemisphere of the unit sphere with minimal angular separation $\pi/3$.¹

L. Szabó (1991) using the Odlyzko - Sloane bound $k(4) \leq 25$ [18] has proved that $B(4) \leq 20$. Recently, K. Bezdek [2, 3] based on $k(4) = 24$ showed that $B(4) \leq 19$ and conjectured that $B(4) = 18$.

We present in this paper a proof that $B(4) = 18$ which is based on our extension of the Delsarte method.

2 On one-sided kissing arrangement

We denote by $\mathbf{S}^{n-1}$ the unit sphere in $\mathbf{R}^n$,

$$\mathbf{S}^{n-1} := \{(x_1, \dots, x_{n-1}, x_n) \in \mathbf{R}^n : x_1^2 + \dots + x_n^2 = 1\}$$

Let S_+ be the closed hemisphere of $\mathbf{S}^{n-1}$ with $x_n \geq 0$, i.e S_+ is a spherical cap of center N and angular radius 90° , where the “North pole” $N = (0, \dots, 0, 1)$. The boundary of S_+ is called the equator and is denoted by E . ($E \cong \mathbf{S}^{n-2}$.)

Let us for $p = (x_1, \dots, x_{n-1}, x_n)$ denote by $p' = (x_1, \dots, x_{n-1}, -x_n)$, and let for $P \subset S_+$ denote by $P' = \{p' : p \in P\}$.

Let $\Pi_n : \dot{S}_+ \rightarrow E$, $\dot{S}_+ := S_+ \setminus \{N\}$, be the projection from the pole N which sends the point $p \in \dot{S}_+$ along its meridian to the equator of the sphere. $Q = \{q_1, \dots, q_m\} \subset E$ is said to be the projection of $P = \{p_1, \dots, p_m\} \subset \dot{S}_+$ if $q_i = \Pi_n(p_i)$. Then $q_i = q_j$ if and only if p_i and p_j lie in the same meridian.

¹Actually, the set $\{p_i\}$ is the part of the vertices of the “24-cell” that belong to S_+ .

Throughout this paper we use the function $\omega(\varphi, \alpha, \beta)$ defined by

$$\cos \omega(\varphi, \alpha, \beta) := \frac{\cos \varphi - \cos \alpha \cos \beta}{\sin \alpha \sin \beta}.$$

This function describes the change of distance φ between two points on $\dot{S}_+$ which are α and β away from N under the action Π_n .

For fixed $\varphi \leq 90^\circ$ and $0 < u \leq \alpha \leq \beta \leq v \leq 90^\circ$, the function $\omega(\varphi, \alpha, \beta)$ attains its minimum at $\alpha = \beta = v$ or $\alpha = u, \beta = v$. If $v < \varphi$, then the minimum of ω achieves at $\alpha = \beta = v$ (see Theorem 3 in [17]).

A set P in $\mathbf{S}^{n-1}$ is called φ -code if the angular separation between any two points from P is at least φ . Denote by $A(n, \varphi)$ the maximal size of φ -codes in $\mathbf{S}^{n-1}$. By $A(n, \varphi, \psi)$ denote the maximal size of φ -codes in a spherical cap with angular radius ψ . So we have $A(n, \varphi) = A(n, \varphi, 180^\circ)$, $k(n) = A(n, 60^\circ)$, $B(n) = A(n, 60^\circ, 90^\circ)$.

We say that a set $P \subset \mathbf{S}^{n-1}$ is a *kissing arrangement* in $\Delta \subset \mathbf{S}^{n-1}$ if $P \subset \Delta$ and P is a 60° -code. P is called a *one-sided kissing arrangement* if $\Delta = S_+$.

Definition. Let $P \subset S_+$. Then

$$P_a := \{p \in P : 90^\circ \geq \text{dist}(p, N) \geq 60^\circ\}, \quad a(P) := |P_a|,$$

$$P_b = \{p \in P : \text{dist}(p, N) < 60^\circ\}, \quad b(P) := |P_b|.$$

So we have

$$P = P_a \cup P_b, \quad |P| = a(P) + b(P).$$

Theorem 1. Let $P \subset \mathbf{S}^{n-1}$ be a one-sided kissing arrangement. Then

$$a(P) + 2b(P) \leq k(n).$$

Proof. For any distinct points $p, q \in P$ we have $\text{dist}(p, q) \geq 60^\circ$. It is easy to see (Fig. 2) that $\text{dist}(p', q) \geq \text{dist}(p, q) \geq 60^\circ$. Note that $\text{dist}(P_b, P'_b) \geq 60^\circ$. Therefore, $\tilde{P} = P \cup P'_b$ is a kissing arrangement in $\mathbf{S}^{n-1}$. Thus, $|\tilde{P}| \leq k(n)$. $\square$




Fig. 2

Theorem 2. Let $P \subset \mathbf{S}^{n-1}$ be a one-sided kissing arrangement. Then

$$a(P) \leq A(n-1, \arccos(1/\sqrt{3})).$$

Proof. Let $Q_a = \{q_i\}$ be the projection of $P_a = \{p_i\}$, $q_i = \Pi_n(p_i)$ (see Fig. 3), and let

$$\theta_i = \text{dist}(N, p_i), \quad \phi_{i,j} = \text{dist}(p_i, p_j), \quad \gamma_{i,j} = \text{dist}(q_i, q_j).$$

Then $\gamma_{i,j} = \omega(\phi_{i,j}, \theta_i, \theta_j)$. We have $\phi_{i,j} \geq 60^\circ$, then $\gamma_{i,j} \geq \omega(60^\circ, \theta_i, \theta_j)$. Since $60^\circ \leq \theta_i \leq 90^\circ$ for all i , we see that

$$\gamma_{i,j} \geq \lambda = \omega(60^\circ, 60^\circ, 90^\circ) = \arccos(1/\sqrt{3}).$$

Then Q_a is a λ -code in $\mathbf{S}^{n-2}$. Thus, $a(P) = |P_a| = |Q_a| \leq A(n-1, \lambda)$. $\square$




Fig. 3

Corollary 1. $([9, 20, 1]) \quad B(3) = 9$.

Proof. Suppose P is a one-sided kissing arrangement in $\mathbf{S}^2$ with $|P| = B(3)$. Since $k(3) = 12$, Theorem 1 implies $a + 2b \leq 12$, where $a = a(P)$, $b = b(P)$. Note that $A(2, \psi) = \lfloor 2\pi/\psi \rfloor$. Therefore, $A(2, \arccos(1/\sqrt{3})) = 6$. Then Theorem 2 yields $a \leq 6$. Also we have $a + b = B(3) \geq 9$. So these nonnegative integer numbers a and b satisfy the following inequalities:

$$a + 2b \leq 12, \quad a \leq 6, \quad a + b \geq 9.$$

It is easy to see that there exists only one solution ($a = 6$, $b = 3$) of these inequalities. Thus, $B(3) = a + b = 9$. $\square$

Corollary 2. *Let $P \subset \mathbf{S}^3$ be a one-sided kissing arrangement with $|P| = B(4)$. Then there are only the following possibilities for $|P|, a(P), b(P)$:*

$$|P| = 19, \quad (a, b) = (15, 4); (14, 5)$$

$$|P| = 18, \quad (a, b) = (15, 3); (14, 4); (13, 5); (12, 6)$$

Proof. Recently we proved that $k(4) = 24$ [15, 17]. That implies $a + 2b \leq 24$. On the other hand, Delsarte's bound gives $A(3, \arccos(1/\sqrt{3})) \leq 15$ (see [5]).² Therefore, we have the following inequalities for integer numbers a, b :

$$a + 2b \leq 24, \quad a \leq 15, \quad a + b = B(4) \geq 18.$$

²For $\lambda = \arccos(1/\sqrt{3}) \approx 54.74^\circ$ Delsarte's bound is stronger than Fejes Tóth's bound [10] $A(3, \lambda) \leq 16$. However, Delsarte's bound also exceeds the lower bound that equals 14. So there is possibility to improve an upper bound for $A(3, \lambda)$ and to skip the case (15,4).

There are only the following integer solutions of these inequalities:
 $(a, b) = (15, 4); (14, 5); (15, 3); (14, 4); (13, 5); (12, 6)$. $\square$

This corollary immediately yields

Corollary 3. (K. Bezdek) $B(4) = 18$ or 19 .

3 An extension of Delsarte's method

For the proof $(a, b) \neq (15, 4), (14, 5)$ we are using techniques that is called “Delsarte's method”. Here we give a brief introduction to this method (for more details see [5, 6, 7, 8, 12, 14, 15, 16, 17, 18, 19]) and define its extension.

Let $P = \{p_1, p_2, \dots, p_M\}$ be any finite subset of $\mathbf{S}^{n-1}$. By $\phi_{i,j}$ we denote the spherical (angular) distance between p_i, p_j , i.e. $\phi_{i,j} := \text{dist}(p_i, p_j)$.

Let us recall the definition of Gegenbauer polynomials by recurrence formula:

$$G_0^{(n)} = 1, \quad G_1^{(n)} = t, \quad G_2^{(n)} = \frac{nt^2 - 1}{n-1}, \quad \dots,$$

$$G_k^{(n)} = \frac{(2k+n-4)t G_{k-1}^{(n)} - (k-1)G_{k-2}^{(n)}}{k+n-3}$$

Note that $G_k^{(n)}(1) = 1$.

Theorem (Schoenberg, 1942). *If $g_{i,j} = G_k^{(n)}(\cos \phi_{i,j})$, then the matrix $(g_{i,j})$ is positive semidefinite.*

The converse holds also: if $f(t)$ is a real polynomial and for any finite $P \subset \mathbf{S}^{n-1}$ the matrix $(f(\cos \phi_{i,j}))$ is positive semidefinite, then f is a sum of $G_k^{(n)}$ with nonnegative coefficients.

The following theorem is a simple consequence of Schoenberg's theorem.

Theorem (Delsarte et al, 1977; Kabatiansky & Levenshtein, 1978). *Assume that a polynomial f satisfies the following property:*

$$f(t) = f_0 G_0^{(n)}(t) + \dots + f_d G_d^{(n)}(t) \quad \text{with} \quad f_0 \geq 0, \dots, f_d \geq 0; \quad (*)$$

Then for any $P = \{p_1, p_2, \dots, p_M\} \subset \mathbf{S}^{n-1}$ we have

$$S_f(P) := \sum_{i=1}^M \sum_{j=1}^M f(\cos \phi_{i,j}) \geq f_0 M^2.$$

Theorem 3. *Let $P = \{p_1, p_2, \dots, p_M\}$, $p_i \in \mathbf{S}^{n-1}$. Suppose f satisfies $(*)$, and $f(t) \leq 0$ for all $t \in [-1, \cos \psi]$. Then*

$$f_0 M^2 \leq f(1)M + T_f(P), \quad \text{where} \quad T_f(P) := \sum_{(i,j): \phi_{i,j} < \psi, i \neq j} f(\cos \phi_{i,j}),$$

Proof. Since $\phi_{i,i} = 0$, we see that $f(\cos \phi_{i,i}) = f(1)$. By the assumption: if $\phi_{i,j} \geq \psi$, then $f(\cos \phi_{i,j}) \leq 0$. Therefore, we have $S_f(P) \leq f(1)M + T_f(P)$. On the other hand, $S_f(P) \geq f_0 M^2$. $\square$

Corollary 4. (Delsarte's bound) *Suppose f satisfies the assumptions of Theorem 3 and $f_0 > 0$. Then*

$$A(n, \varphi) \leq \frac{f(1)}{f_0}.$$

Proof. Let P be a φ -code of size M on $\mathbf{S}^{n-1}$, i.e. $\phi_{i,j} \geq \varphi$ for $i \neq j$. Then $T_f(P) = 0$. Thus, Theorem 3 yields $f_0 M^2 \leq f(1)M$. $\square$

Remark. Suitable polynomials f can be found by the linear programming method. See details and algorithms in [6, 7, 8, 12, 17, 18].

Now we introduce some extensions of Delsarte's method for one-sided kissing arrangements. Note that all results in this section hold for much more general case: φ -codes in spherical caps. However, for simplicity, we consider here only one-sided kissing arrangements.

Let $P \subset S_+ \subset \mathbf{S}^{n-1}$ be a one-sided kissing arrangement. For fixed θ_0 , where $60^\circ < \theta_0 < 90^\circ$, we define two subsets of P_a :

$$P_{a_1} = \{p_i \in P_a : \theta_0 < \text{dist}(p_i, N)\}, \quad P_{a_2} = \{p_i \in P_a : \theta_0 \geq \text{dist}(p_i, N)\},$$

$$a_1 = a_1(P) := |P_{a_1}|, \quad a_2 = a_2(P) := |P_{a_2}|, \quad a = a_1 + a_2.$$

Recall that $|P| = a + b = a_1 + a_2 + b$.

Theorem 4. *Suppose a polynomial f satisfies $(*)$, $f(t) \leq 0$ for all $t \in [-1, 1/2]$, and $f(t)$ is a monotone increasing function on $[1/2, 1]$. Then*

$$(a_1 + 2a_2 + 2b)^2 f_0 \leq (a_1 + 2a_2 + 2b)f(1) + 2a_2 f(-\cos 2\theta_0).$$

Proof. Let

$$\tilde{P} = P \cup P_{a_2} \cup P'_b.$$

Then $M = |\tilde{P}| = |P| + a_2 + b = a_1 + 2a_2 + 2b$.

If $p \in P_{a_2}$, then $\text{dist}(p, p') \geq 180^\circ - 2\theta_0$. For any other pair (p, q) from $\tilde{P}$ we have $\text{dist}(p, q) \geq 60^\circ$ (see Theorem 1, Fig. 2). Therefore, the monotonicity assumption implies: $T_f(\tilde{P}) \leq 2a_2 f(\cos(180^\circ - 2\theta_0)) = 2a_2 f(-\cos 2\theta_0)$. Thus, the proof follows from Theorem 3 with $\psi = 60^\circ$. $\square$

Now for any function $f(t)$ and θ_0 , $60^\circ < \theta_0 < 90^\circ$, we introduce the function $R_f(\theta)$ on the interval $[0, \theta_0]$.

Let

$$\psi_0 := \omega(60^\circ, \theta_0, 90^\circ) = \arccos\left(\frac{1}{2 \sin \theta_0}\right).$$

For given $e_0 \in E \cong \mathbf{S}^{n-2}$ denote by $C(e_0, \psi_0)$ a spherical cap in the equator E of center e_0 and radius ψ_0 . Let $\Lambda_n(\theta_0)$ be a spherical cone in $\mathbf{S}^{n-1}$ with vertex

N and with the base $C(e_0, \psi_0)$ (Fig. 4). In other words, $\Lambda_n(\theta_0)$ is a cone of height 90° and radius ψ_0 .

Consider a point p in the height Ne_0 with $\text{dist}(N, p) = \theta \leq \theta_0$. Denote by $\Delta_n(\theta, \theta_0)$ the domain in $\Lambda_n(\theta_0)$ that lies below the sphere $S(p) = S(p, 60^\circ)$ of center p and radius 60° . It is easy to see that for $\theta \leq \theta_0$ this domain is nonempty.

Definition. Consider kissing arrangements $X = \{p_1, \dots, p_m\}$ in $\Delta_n(\theta, \theta_0)$. Let $\mu(n, \theta, \theta_0)$ be the highest value of m . By $\Omega_n(\theta, \theta_0)$ we denote the set of all kissing arrangements in $\Delta_n(\theta, \theta_0)$.




Fig. 4

Let $Y = \{q_1, \dots, q_m\}$ be the projection of $X = \{p_1, \dots, p_m\} \subset \Delta_n(\theta, \theta_0)$, where $q_i = \Pi_n(p_i)$, and let

$$F_f(X) = F_f(p_1, \dots, p_m) := f(\cos \varphi_1) + \dots + f(\cos \varphi_m), \quad \varphi_i := \text{dist}(e_0, q_i).$$

We define $R_f(\theta; \theta_0, n)$ as the maximum of $F_f(X)$ on $\Omega_n(\theta, \theta_0)$:

$$R_f(\theta) = R_f(\theta; \theta_0, n) := \sup_{X \in \Omega_n(\theta, \theta_0)} \{F_f(X)\}.$$

Suppose $P = \{p_1, \dots, p_\ell\} \subset \mathbf{S}^{n-1}$ lies in the cap of center N and radius θ_0 . Let

$$\tilde{R}_f(P) = \tilde{R}_f(P; \theta_0, n) := R_f(\theta_1) + \dots + R_f(\theta_\ell), \quad \theta_k := \text{dist}(N, p_k).$$

Proposition. $R_f(\theta)$ is a monotone decreasing function in θ .

Proof. For $\alpha > \beta$, we have $\Delta_n(\alpha, \theta_0) \subset \Delta_n(\beta, \theta_0)$, then $\Omega_n(\alpha, \theta_0) \subset \Omega_n(\beta, \theta_0)$, so then $R_f(\alpha) \leq R_f(\beta)$. $\square$

Theorem 5. Let $\theta_0 \in (60^\circ, 90^\circ)$. Suppose a polynomial f satisfies the following assumptions: $f = f_0 G_0^{(n-1)} + \dots + f_d G_d^{(n-1)}$ with $f_0 \geq 0, \dots, f_d \geq 0$; and $f(t) \leq 0$ for all $t \in [-1, \cos \psi_0]$.

Then for any one-sided kissing arrangement $P \subset \mathbf{S}^{n-1}$ with $|P| = M$ we have

$$f_0 M^2 \leq M f(1) + 2a_2 R_f(60^\circ; \theta_0, n) + 2\tilde{R}_f(P; \theta_0, n).$$

Proof. First let us prove the theorem for the case when the North pole N is not a point in P . In this case the projection Q of P is well defined.

If we apply Theorem 3 for f and $Q = \Pi_n(P) \subset E$, then we obtain

$$f_0 M^2 \leq S_f(Q) \leq f(1)M + T_f(Q).$$

From the assumption: $f(t) \leq 0$, $t \in [-1, \cos \psi_0]$, for $q_i, q_j \in Q$ it follows that if $\text{dist}(q_i, q_j) \geq \psi_0$, then $f(\cos \phi_{i,j}) \leq 0$. Arguing as in Theorem 2 it is easy to see that for any distinct $q_1, q_2 \in Q_{a_1}$ we have $\text{dist}(q_1, q_2) \geq \psi_0$.

For $q_1, q_2 \in Q_{a_2}$ we have

$$\text{dist}(q_1, q_2) \geq \omega(60^\circ, \theta_0, \theta_0) > \omega(60^\circ, \theta_0, 90^\circ) = \psi_0.$$

In the case $q_1, q_2 \in Q_b$ we have

$$\text{dist}(q_1, q_2) > \omega(60^\circ, 60^\circ, 60^\circ) > \psi_0.$$

Therefore, $\text{dist}(q_1, q_2) < \psi_0$ only if

$$(i) \ q_1 \in Q_{a_1}, \ q_2 \in Q_{a_2}, \text{ or } (ii) \ q_1 \in Q_a, \ q_2 \in Q_b.$$

Let $p \in P_{a_2}$, $\text{dist}(N, p) = \theta$, $e_0 = \Pi_n(p)$, and let $X = \{p_1, \dots, p_m\} \subset P$ with $\text{dist}(e_0, q_i) < \psi_0$. By (i) we have $X \subset P_{a_1}$. This implies that X lies below $S(p, 60^\circ)$. Thus, $X \subset \Delta_n(\theta, \theta_0)$ and $F_f(X) \leq R_f(\theta)$. The same holds for $p \in P_b$ (the case (ii)). We obtain

$$T_f(Q) \leq 2\tilde{R}_f(P_{a_2}) + 2\tilde{R}_f(P_b).$$

By definition $\text{dist}(N, p) \geq 60^\circ$ for $p \in P_{a_2}$. Since $R_f(\theta)$ is a monotone decreasing function, we see that $\tilde{R}_f(P_{a_2}) \leq a_2 R_f(60^\circ)$.

It can be shown in the usual way that the theorem for $N \notin P$ implies the theorem for $N \in P$. Indeed, if we shift p to some point $p(\varepsilon)$ which is a small distance ε away from N , then we can apply the theorem for a φ -code $P(\varepsilon) = p(\varepsilon) \cup P \setminus p$, where $\varphi = 60^\circ - \varepsilon$. Since $R_f(\theta)$ is a continuous function and $R_f(0)$ is well defined the theorem for $N \in P$ follows whenever $\varepsilon \rightarrow 0$. $\square$

In the end of this section we consider some geometric properties of kissing arrangements in $\Delta_n(\theta, \theta_0)$. These properties will help us to compute $R_f(\theta)$ in the next section.

Note that for $\theta > 30^\circ$ the sphere $S(p, 60^\circ)$ intersects the equator. That yields

$$\text{dist}(e_0, q_i) \geq \rho(\theta) := \omega(60^\circ, \theta, 90^\circ) = \arccos\left(\frac{1}{2\sin\theta}\right), \quad \theta > 30^\circ$$

It is easy to see that $p_i \in \Delta_n(\theta, \theta_0)$ implies $\text{dist}(N, p_i) \geq \gamma(\theta, \theta_0)$, where the function $\gamma = \gamma(\theta, \theta_0) = \gamma_1(\theta, 60^\circ, \psi_0)$ can be defined by the equation

$$\omega(60^\circ, \theta, \gamma) = \psi_0, \quad \gamma \geq \theta.$$

In other words, $\gamma_1(\theta, \varphi, \psi)$ is the length of AB of a spherical triangle ABC with $|AC| = \theta$, $|BC| = \varphi$, and $\angle BAC = \psi$. In fact, this equation has two solutions. Let us denote the second solution by $\gamma_2(\theta, \varphi, \psi)$.

We get

$$\text{dist}(q_i, q_j) \geq d(\theta) = d(\theta, \theta_0) := \omega(60^\circ, \gamma(\theta, \theta_0), 90^\circ), \quad i \neq j.$$

Therefore, the set $Y = \{q_i\}$ is a $d(\theta)$ -code on the cap $C(e_0, \psi_0) \subset E \cong \mathbf{S}^{n-2}$. Thus,

$$\mu(n, \theta, \theta_0) \leq A(n-1, d(\theta, \theta_0), \psi_0).$$

It was proved (see Theorem 3 in [17]) that for $\varphi > \psi$ we have

$$A(n, \varphi, \psi) = A(n-1, \omega(\varphi, \psi, \psi)).$$

By $\theta_*(\theta_0)$ denote the solution of the equation $d(\theta, \theta_0) = \psi_0$. We obtain

$$\mu(n, \theta, \theta_0) \leq A(n-2, \omega(d(\theta), \psi_0, \psi_0)) \quad \text{for } \theta > \theta_*(\theta_0).$$

Consider the case $n = 4$. We have a $d(\theta)$ -code $Y = \{q_1, \dots, q_m\}$ on the cap $C(e_0, \psi_0) \subset \mathbf{S}^2$. Now we consider two cases: a) $\theta > \theta_*(\theta_0)$ and b) $\theta \leq \theta_*(\theta_0)$.

a) By definition for $\theta > \theta_*(\theta_0)$ we have $d(\theta) > \psi_0$. It can be shown (see Corollary 2 in [17]) that in this case $m \leq 5$.

Let $m = 5$. Then q_i are vertices of a spherical convex pentagon [17, Theorem 2]. We denote this pentagon by V_5 . Since $\text{dist}(q_i, q_j) \geq d(\theta)$ we have that the length of any side of V_5 is at least $d(\theta)$. On the other hand, $V_5 \subset C(e_0, \psi_0)$. It is not hard to prove that the minimum distance between e_0 and a vertex q_i of V_5 achieves when V_5 is an equilateral pentagon of sides lengths $d(\theta)$ and four other vertices q_j lie in the boundary of the cap $C(e_0, \psi_0)$. That yields

$$\text{dist}(e_0, q_i) \geq \rho_5(\theta, \theta_0) := \gamma_2(\psi_0, d(\theta), \eta), \quad \eta = 180^\circ - 3\omega(d(\theta), \psi_0, \psi_0)/2.$$

b) In this case $m \leq 6$. The maximal size arrangement can be achieved when $q_6 = e_0$ and $\{q_1, \dots, q_5\}$ are vertices of a spherical convex pentagon.

Indeed, let $\delta = \omega(d(\theta), \psi_0, \psi_0)$. It can be shown numerically that for any $\theta_0 \in (60^\circ, 90^\circ)$ we have $\delta > 60^\circ$. Let V be the (spherical) convex hull of Y . It can be shown that $\angle q_i e_0 q_j \geq \delta$, where q_i, q_j are vertices of V . Since $\lfloor 360^\circ/\delta \rfloor \leq 5$ we have that the number of vertices of V is at most 5. It's not hard to prove that no more than one $q_i \in Y$ can lie inside V . Thus, $m \leq 6$.

These facts can be summarized as follows:

Theorem 6. Suppose $X = \{p_1, \dots, p_m\}$ is a kissing arrangement in $\Delta_n(\theta, \theta_0)$. Let $q_i = \Pi_n(p_i)$. Then

- 1) $\text{dist}(q_i, q_j) \geq d(\theta, \theta_0)$, and $\text{dist}(e_0, q_i) \geq \rho(\theta)$ for $\theta > 30^\circ$;
- 2) $m \leq \mu(n, \theta, \theta_0) \leq A(n-1, d(\theta, \theta_0), \psi_0)$;
- 3) $m \leq \mu(n, \theta, \theta_0) \leq A(n-2, \omega(d(\theta, \theta_0), \psi_0, \psi_0))$ for $\theta > \theta_*(\theta_0)$;
- 4) if $n = 4$, then $m \leq \mu(4, \theta, \theta_0) \leq 6$; moreover for $\theta > \theta_*(\theta_0)$ we have $m \leq 5$, and if $m = 5$, then $\text{dist}(e_0, q_i) \geq \rho_5(\theta, \theta_0)$.

4 B(4)=18

In this section we prove the main theorem of this paper:

Theorem 7. $B(4) = 18$.

Now we prove that $(a, b) \neq (15, 4), (14, 5)$. In this section $\theta_0 = 71.74^\circ$. So we have $\psi_0 \approx 58.2299^\circ$, $\theta_* = \theta_*(\theta_0) \approx 25.8526^\circ$, and $\cos 2\theta_0 = -\cos 36.52^\circ$. Here and below numbers are shown to four decimal places.

Lemma 1. Let $P \subset \mathbf{S}^3$ be a one-sided kissing arrangement with $|P| = 19$.

- 1) If $(a, b) = (15, 4)$, then $a_2 \leq 5$.
- 2) If $(a, b) = (14, 5)$, then $a_2 \leq 3$.

Proof. Let $f_{OS}(t) = f_0 + f_1 G_1^{(4)}(t) + \dots + f_9 G_9^{(4)}(t)$, where $f_0 = 1$, $f_1 = 3.6181$, $f_2 = 6.1156$, $f_3 = 7.0393$, $f_4 = 5.0199$, $f_5 = 2.313$, $f_6 = f_7 = f_8 = 0$, $f_9 = 0.4525$.

The polynomial f_{OS} satisfies the assumptions of Theorem 4.³ Now we apply this theorem for the proof.

- 1) Assume the converse. Then $a_2 \geq 6$, and $M = a_1 + 2a_2 + 2b = |P| + a_2 + b \geq 29$. For $a_2 = 6$ from Theorem 4

$$841 = f_0 29^2 \leq 29 f_{OS}(1) + 12 f_{OS}(\cos(36.52^\circ)) \approx 840.8819$$

follows, a contradiction.

- 2) If $(a, b) = (14, 5)$ and $a_2 = 4$, then $M = 28$. We have

$$784 = f_0 28^2 \leq 28 f_{OS}(1) + 8 f_{OS}(\cos(36.52^\circ)) \approx 782.0941,$$

a contradiction.

Since $f_0 M^2 - M f_{OS}(1) - 2a_2 f_{OS}(\cos(36.52^\circ))$ is increasing whenever M is increasing, we complete our proof. $\square$

Lemma 2. $(a, b) \neq (15, 4)$.

Proof. Assume the converse.

Let $g(t) = g_0 + g_1 G_1^{(3)}(t) + \dots + g_9 G_9^{(3)}(t)$, where $g_0 = 1$, $g_1 = 2.7986$, $g_2 = 3.6388$, $g_3 = 3.4429$, $g_4 = 2.1227$, $g_5 = 0.8637$, $g_6 = g_7 = g_8 = 0$, $g_9 = 0.1281$. Then $g(1) = g_0 + g_1 + \dots + g_9 = 13.9948$.

The polynomial g satisfies the assumptions of Theorem 5 with $n = 4$. Note that $g(t)$ is a monotone increasing function on $[\cos \psi_0, 1]$.

Let us apply Theorem 5 for P_a . Since $|P_a| = 15$ and $b(P_a) = 0$ we have

$$225 = g_0 15^2 \leq 15 g(1) + 2a_2 R_g(60^\circ).$$

Now we prove that $R_g(60^\circ) \leq 4g(1/\sqrt{3})$.

³This polynomial was applied by Odlyzko and Sloane [18] to prove that $k(4) \leq 25$. Indeed, Delsarte's bound gives $k(4) = A(4, 60^\circ) \leq f_{OS}(1)/f_0 = f_{OS}(1) \approx 25.5584$. Then $k(4) \leq 25$.

Theorem 6 yields

$$\text{dist}(e_0, q_i) \geq \rho(60^\circ) = \arccos(1/\sqrt{3}) \approx 54.7356^\circ,$$

$\mu(4, 60^\circ, \theta_0) \leq 5$, and for $m = 5$ we have

$$\text{dist}(e_0, q_i) \geq \rho_5 = \rho_5(60^\circ, \theta_0) \approx 57.5043^\circ.$$

Then the monotonicity of $g(t)$ on $[\cos \psi_0, 1]$ implies for $m = 5$: $F_g(X) \leq 5g(\cos \rho_5)$ and for $m \leq 4$: $F_g(X) \leq 4g(1/\sqrt{3})$. Since

$$4g(1/\sqrt{3}) \approx 1.4673 > 5g(\cos \rho_5) \approx 0.332,$$

we have $R_g(60^\circ) \leq 4g(1/\sqrt{3})$. (It is easy to see that $R_g(60^\circ) = 4g(1/\sqrt{3})$.)

By Lemma 1, $a_2 \leq 5$, so that

$$225 \leq 15g(1) + 2a_2R_g(60^\circ) \leq 15g(1) + 40g(1/\sqrt{3}) \approx 224.5946$$

- a contradiction. □

Lemma 3. $(a, b) \neq (14, 5)$.⁴

Proof. Assume the converse.

1. Let $P_b = \{p_1, \dots, p_5\}$, $\theta_i := \text{dist}(p_i, N)$, and let $\{\bar{p}_i\}$ be sorted $\{p_i\}$ by θ_i :

$$\{p_i\} = \{\bar{p}_j\}, \quad \bar{\theta}_i := \text{dist}(\bar{p}_i, N), \quad 60^\circ > \bar{\theta}_1 \geq \bar{\theta}_2 \geq \bar{\theta}_3 \geq \bar{\theta}_4 \geq \bar{\theta}_5 > 0.$$

By $P_b(k)$ denote $\{\bar{p}_1, \dots, \bar{p}_k\}$. Here $1 \leq k \leq 5$, and $P_b(5) = P_b$. Let

$$P(k) := P_a \bigcup P_b(k), \quad Q(k) := Q_a \bigcup Q_b(k).$$

Then $|P(k)| = |Q(k)| = 14 + k$.

Suppose a polynomial f satisfies the assumptions of Theorem 5 with $n = 4$.

Then Theorem 5 for $P(k)$ yields

$$f_0(14 + k)^2 \leq S_f(Q(k)) \leq (14 + k)f(1) + 2a_2R_f(60^\circ) + 2\tilde{R}_f(P_b(k)). \quad (1)$$

Let $h(t) = h_0 + h_1G_1^{(3)}(t) + \dots + h_9G_9^{(3)}(t)$, where
 $h_0 = 1$, $h_1 = 2.722$, $h_2 = 3.6191$, $h_3 = 3.4323$, $h_4 = 2.4227$, $h_5 = 1.1991$,
 $h_6 = h_7 = h_8 = 0$, $h_9 = 0.0195$. $h(1) = h_0 + h_1 + \dots + h_9 = 14.7413$.

Note that h (as well as g) satisfies the assumptions of Theorem 5, and $h(t)$ is a monotone increasing function on $[\cos \psi_0, 1]$. Actually, g and h are close to each other. As above, $R_h(60^\circ) = 4h(1/\sqrt{3})$. However, for $k \leq 3$ using $f = g$ in (1) we get better results than for $f = h$, but h is better than g for $k = 4, 5$.

⁴There is old conjecture that the maximal kissing arrangement in four dimensions is unique up to isometry, i.e. is the “24-cell” P_{24} . From this conjecture easily follows Lemma 3. Indeed, let $\hat{P} := P \bigcup P'_b$, then $|\hat{P}| = a + 2b = 24$. The conjecture yields that $\text{conv}(\hat{P})$ is the “24-cell”. Since at most 18 vertices of P_{24} can lie in a closed hemisphere, $|\hat{P}| \leq 18$ - a contradiction.

By Lemma 1 $a_2 \leq 3$. Therefore, for $f = g$ and $f = h$ we have

$$(14 + k)^2 \leq (14 + k)f(1) + 24f(1/\sqrt{3}) + 2\tilde{R}_f(P_b(k)). \quad (2)$$

Recall that $\tilde{R}_f(P_b(k)) = R_f(\bar{\theta}_1) + \dots + R_f(\bar{\theta}_k)$. Thus, (2) yields

$$R_f(\bar{\theta}_1) + \dots + R_f(\bar{\theta}_k) \geq k^2/2 + 14k + 98 - (7 + k/2)f(1) - 12f(1/\sqrt{3}). \quad (3)$$

2. Now we compute the upper bound on $R_f(\theta)$ for $f = g, h$, and $0 \leq \theta \leq 60^\circ$.

Let $\theta \in [0^\circ, 60^\circ]$. If X is a kissing arrangement in $\Delta_4(\theta, \theta_0)$, then Theorem 6 yields some constraints for Y , where Y is the projection of X . Using these constraints we introduce the function $L_f(\theta)$ such that $R_f(\theta) \leq L_f(\theta)$.

For a fixed $e_0 \in \mathbf{S}^2$ we consider the set $\Psi(\theta)$ of all $d(\theta)$ -codes $Y = \{q_1, \dots, q_m\}$ on $\mathbf{S}^2$ such that $m \leq 6$; $\varphi_i := \text{dist}(e_0, q_i) \geq \rho(\theta)$ for $\theta > 30^\circ$; $m \leq 5$ for $\theta > \theta_*$ and if $m = 5$ then $\varphi_i \geq \rho_5(\theta)$. Let

$$H_f(Y) := \tilde{f}(\varphi_1) + \dots + \tilde{f}(\varphi_m), \quad L_f(\theta) := \sup_{Y \in \Psi(\theta)} \{H_f(Y)\},$$

where

$$\tilde{f}(\varphi) := \begin{cases} f(\cos \varphi) & 0 \leq \varphi \leq \psi_0 \\ 0 & \varphi > \psi_0 \end{cases}$$

Theorem 6 implies that if $X \in \Omega_4(\theta, \theta_0)$, then $Y \in \Psi(\theta)$. Thus $R_f(\theta) \leq L_f(\theta)$.

Consider some $\theta \in [30^\circ, 60^\circ]$. Recall that $f(\cos \varphi)$ is a monotone decreasing function on $[0, \psi_0]$. Then (as well as in Lemma 2) we have for $m = 5$: $H_f(Y) \leq 5\tilde{f}(\rho_5(\theta))$, and for $m \leq 4$: $H_f(Y) \leq 4\tilde{f}(\rho(\theta))$. It can be shown numerically that in this interval $4\tilde{f}(\rho(\theta)) > 5\tilde{f}(\rho_5(\theta))$. Therefore, $L_f(\theta) \leq 4\tilde{f}(\rho(\theta))$.

Let $\delta_4 \approx 44.4577^\circ$ is defined by the equality: $\cos^2 \rho(\delta_4) = \cos d(\delta_4)$. This equality means that a spherical square of side length $d(\delta_4)$ has the circumradius equals $\rho(\delta_4)$. Then for $\theta \in [\delta_4, 60^\circ]$ the side length of a spherical square with the circumcircle of center e_0 and radius $\rho(\theta)$ is not less than $d(\theta)$. Thus,

$$L_f(\theta) = 4\tilde{f}(\rho(\theta)) = 4f\left(\frac{1}{2\sin \theta}\right) \quad \text{for } \theta \in [\delta_4, 60^\circ].$$

If $\theta < \delta_4$, then the circumradius of a square of side length $d(\theta)$ is greater than $\rho(\theta)$. It have been proved (see Theorem 4 [17]) that if Y is optimal (i.e. $H_f(Y) = L_f(\theta)$) and $|Y| = 4$, then Y is a spherical rhomb $q_1q_2q_3q_4$ with edge lengths $d(\theta)$.

Consider a rhomb $Y = q_1q_2q_3q_4$ of side lengths $d(\theta)$ with $\text{dist}(q_1, q_3) \geq \text{dist}(q_2, q_4)$ such that the circumradius of $q_1q_2q_4$ is $\rho(\theta)$. Under these conditions this rhomb there exists and uniquely defined for $\theta \in [\delta_3, \delta_4]$. Here $\delta_3 \approx 37.4367^\circ$ is defined by the condition that $\rho(\delta_3)$ is the circumradius of a regular triangle of side lengths $d(\delta_3)$. It can be verified numerically that for all $\theta \in [\delta_3, \delta_4]$ $H_f(Y)$ achieves its maximum when e_0 is the circumcenter of $q_1q_2q_4$. Since for this arrangement $\text{dist}(e_0, q_3)$ is uniquely defined by θ we denote it by $r_4(\theta)$. Then

$$L_f(\theta) = 3f\left(\frac{1}{2\sin \theta}\right) + \tilde{f}(r_4(\theta)) \quad \text{for } \theta \in [\delta_3, \delta_4].$$

Note that for $\theta = \delta_3$ we have $\varphi_3 > \psi_0$, i.e. $\tilde{f}(\varphi_3) = 0$. Computations show that for $\theta \in [30^\circ, \delta_3]$ the maximum of $H_f(Y)$ is achieved when $q_1 q_2 q_4$ is a regular triangle of side lengths $d(\theta)$ with $\text{dist}(e_0, q_1) = \rho(\theta)$. Let $r_3(\theta) := \text{dist}(e_0, q_2) = \text{dist}(e_0, q_4)$. Since in this case $\varphi_3 > \psi_0$, we obtain

$$L_f(\theta) = f\left(\frac{1}{2\sin\theta}\right) + 2\tilde{f}(r_3(\theta)) \quad \text{for } \theta \in [30^\circ, \delta_3].$$

For $\theta = 30^\circ$ we see that $q_1 = e_0$ and $r_3(\theta) = d(\theta) > \psi_0$. Therefore, $L_f(30^\circ) = f(1)$. Note that for $\theta \in [30^\circ, \theta_*]$ we have $f(1) > 5\tilde{f}(\rho_5(\theta))$. Then, as above, we can assume that Y is a rhomb. Computations show that $H_f(Y)$ attains its maximum when e_0 is a vertex of Y , i.e. $L_f(\theta) = f(1)$. For $\theta < \theta_*$ it is not hard to show that $H_f(Y)$ achieves its maximum when $e_0 \in Y$ and other q_i lie at the distance $d(\theta)$ away from e_0 . Thus,

$$L_f(\theta) = f(1) + 5\tilde{f}(d(\theta)) \quad \text{for } \theta \in [0, 30^\circ].$$

(Note that if $\theta \geq \theta_*$, then $d(\theta) \geq \psi_0$, i.e. $\tilde{f}(d(\theta)) = 0$.)

3. Consider the domain $\Theta = \{(\theta_1, \dots, \theta_5)\}$ in $\mathbf{R}^5$. The proof of Lemma 3 follows from the fact that Θ is empty set under the constraints (3) and $\text{dist}(p_i, p_j) \geq 60^\circ$.

It is not so hard to find bounds for θ_k under the constraints (3). For instance, since $R_f(\theta)$ is a monotone decreasing function, we have

$$R_f(\bar{\theta}_1) + \dots + R_f(\bar{\theta}_k) \leq kR_f(\bar{\theta}_k) \leq kL_f(\bar{\theta}_k).$$

Combining this with (3) we get

$$\bar{\theta}_1 < 53.93^\circ, \bar{\theta}_2 < 48.45^\circ, \bar{\theta}_3 < 46.72^\circ, \bar{\theta}_4 < 45.53^\circ, \bar{\theta}_5 < 44.47^\circ.$$

Here for $k = 1, 2, 3$ we applied $f = g$, and for $k = 4, 5$: $f = h$.

Moreover, using convexity of $L_f(\theta)$ on $[\delta_4, 60^\circ]$ we can give inequalities for sums of θ_k . Perhaps, it is possible using these inequalities to prove the lemma without computer works. However, we think that is a hard problem. Here for the proof we apply the numerical method that was developed in our paper [17].

Denote by $P_5(\alpha)$ a convex polytope in $\mathbf{S}^3$ with five vertices (see Fig. 5). The lengths of all edges $P_5(\alpha)$, except $p_2 p_4$, $p_3 p_5$, are equal to 60° , and

$$\text{dist}(p_2, p_4) = \alpha \in [60^\circ, 90^\circ], \text{dist}(p_3, p_5) = \beta, \quad 2\cos\alpha\cos\beta + \cos\alpha + \cos\beta = 0.$$




Fig. 5: $P_5(\alpha)$

Let $X = \{p_1, \dots, p_5\}$ be a kissing arrangement in the cap $C(N, 60^\circ) \subset \mathbf{S}^3$, and let

$$F(X) = F_1(\theta_1) + \dots + F_5(\theta_5), \quad \theta_i = \text{dist}(N, p_i),$$

where $F_i(\theta)$ is a monotone decreasing function in θ . From Theorem 5 [17] it follows that F attains its maximum on $\{X\}$ at X_0 when $\text{conv}(X_0)$ (the convex hull of X_0) is isometric to $P_5(\alpha)$ for some $\alpha \in [60^\circ, 90^\circ]$. Note that Theorem 5 is not be assumed a strong monotonicity of $F_i(\theta)$.

Let us apply this fact with $F(P_b) = L_f(\bar{\theta}_1) + \dots + L_f(\bar{\theta}_k)$, i.e. $F_i = L_f$ for $i = 1, \dots, k$, and $F_i = 0$ for $i > k$. (It's easy to see that $L_f(\theta)$ is a monotone decreasing function.) Thus, we can assume that $\text{conv}(P_b) = P_5(\alpha)$.

Let us show how using this fact to obtain a contradiction for the lemma.

Let the vertices p_1, p_2, p_3 of $P_5(\alpha)$ be fixed. Then the vertices $p_4 = p_4(\alpha)$ and $p_5 = p_5(\alpha)$ are uniquely determined by α . Consider the following domain

$$D_\varepsilon(u_1, u_2, u_3) := \{p \in S^3 : \theta_i = \text{dist}(p, p_i) \in [u_i, u_i + \varepsilon]\}, \quad \varepsilon > 0.$$

In general position $D = D_\varepsilon(u_1, u_2, u_3)$ looks like a curvilinear parallelepiped, i.e. D has (at most) eight vertices.

Let us denote by $S = S(u_1, u_2, u_3; \alpha) = \{s_i\}$ the set that consists of vertices of D and points on the boundary of D which are intersections of the great circles through $p_i p_j$, $i \neq j$, $j = 4, 5$, and D . For small ε and $u_i > 0$ it is easy to see that $|S| \leq 10$. Lemma 5 [17] yields that the function $\theta_i = \text{dist}(p, p_i)$, $i = 4, 5$, in p achieves its minimum on D at some s_j . Let us denote this point by $\tilde{p}_i(\alpha, D)$. Thus, $L_f(\theta_i)$ (as well as any monotone decreasing function in θ) achieves its maximum on D at $\theta_i = \tilde{\theta}_i(\alpha) := \text{dist}(p_i, \tilde{p}_i(\alpha, D))$.

Note that θ_4 is increasing and θ_5 is decreasing whenever α is increasing. Then for $\alpha \in [r, r + \varepsilon]$ we have $\tilde{\theta}_4(\alpha) \geq \tilde{\theta}_4(r)$ and $\tilde{\theta}_5(\alpha) \leq \tilde{\theta}_5(r + \varepsilon)$. Finally, for $L_f(\theta_i)$ on $\bar{D}_\varepsilon = D \times [r, r + \varepsilon]$ we have

$$L_f(\theta_i) \leq L_f(u_i), \quad i = 1, 2, 3; \quad L_f(\theta_4) \leq L_f(\tilde{\theta}_4(r)); \quad L_f(\theta_5) \leq L_f(\tilde{\theta}_5(r + \varepsilon)) \quad (4).$$

Using (4) we can easily to find an upper bound on $L_f(\theta_i)$ on any $\bar{D}_\varepsilon$. Then we can check (3) for points in $\bar{D}_\varepsilon$. So if we cover $S_+ \times [60^\circ, 90^\circ]$ by $\bar{D}_\varepsilon$ we can check (3) for all possible p , $\{p_i\}$ and α . Our computations show that there no points in Θ that satisfy (3) - a contradiction. $\square$

5 Conclusions

It is clear that between kissing numbers and one-sided kissing numbers there are some relations. Look at these nice equalities:

$$\begin{aligned} n = 2, \quad 4 = B(2) &= \frac{k(1) + k(2)}{2} = \frac{2 + 6}{2}; \\ n = 3, \quad 9 = B(3) &= \frac{k(2) + k(3)}{2} = \frac{6 + 12}{2}; \end{aligned}$$

$$n = 4, \quad 18 = B(4) = \frac{k(3) + k(4)}{2} = \frac{12 + 24}{2}.$$

We are not expect that the same holds for all dimensions, i.e.

$$B(n) = \bar{K}(n) := \frac{k(n-1) + k(n)}{2}$$

for all n . However, there are many reasons that $B(n) = \bar{K}(n)$ for $n = 5, 8, 24$; $B(n) \approx \bar{K}(n)$ for all n ; and this equality holds asymptotically

$$\lim_{n \rightarrow \infty} \frac{B(n)}{\bar{K}(n)} = 1.$$

If we consider the minimal vectors of D_5, E_8 , and Leech lattices, then we see that $B(5) \geq 32$, $B(8) \geq 183$, $B(24) \geq 144855$. We propose the following

Conjecture. $B(5) = 32$, $B(8) = 183$, $B(24) = 144855$.

Note that our method could be applied to higher dimensions. However, it does not give sharp upper bound on $B(n)$. It's an interesting problem to find better method, in particular for $n = 5, 8, 24$.

Acknowledgment. I wish to thank Károly Bezdek and Gábor Fejes Tóth for helpful discussions and useful comments on this paper.

References

- [1] A. Bezdek and K. Bezdek, A note on the ten-neighbour packing of equal balls, *Beiträge zur Alg. und Geom.* **27** (1988), 49-53.
- [2] K. Bezdek, Sphere packings in 3-space, *Proceedings of the COE Workshop on Sphere Packings*, Kyushu University Press (2004), 32-49.
- [3] K. Bezdek, Sphere packing revisited, *European J. Comb.* **27** (2006), 864-883.
- [4] K. Bezdek and P. Brass, On k^+ -neighbour packings and one-sided Hadwiger configurations, *Contributions to Algebra and Geometry*, **4** (2003), 493-498.
- [5] P.G. Boyvalenkov, D.P. Danev and S.P. Bumova, Upper bounds on the minimum distance of spherical codes, *IEEE Trans. Inform. Theory*, **42**(5), 1996, 1576-1581.
- [6] J.H. Conway and N.J.A. Sloane, *Sphere Packings, Lattices, and Groups*, New York, Springer-Verlag, 1999 (Third Edition).

- [7] Ph. Delsarte, Bounds for unrestricted codes by linear programming, Philips Res. Rep., **27**, 1972, 272-289.
- [8] Ph. Delsarte, J.M. Goethals and J.J. Seidel, Spherical codes and designs, Geom. Dedic., **6**, 1977, 363-388.
- [9] G. Fejes Tóth, Ten-neighbor packing of equal balls, Periodica Math. Hungar., **12** (1981), 125-127.
- [10] L. Fejes Tóth, Lagerungen in der Ebene, auf der Kugel und in Raum, Springer-Verlag, 1953; Russian translation, Moscow, 1958.
- [11] L. Fejes Tóth and H. Sachs, Research problem 17, Periodica Math. Hungar. **7** (1976), 125-127.
- [12] G.A. Kabatiansky and V.I. Levenshtein, Bounds for packings on a sphere and in space, Problems of Information Transmission, **14**(1), 1978, 1-17.
- [13] G. Kertész, Nine points on the hemisphere, Colloq. Math. Soc. J. Bolyai (Intuitive Geometry, Szeged 1991), **63** (1994), 189-196.
- [14] V.I. Levenshtein, On bounds for packing in n -dimensional Euclidean space, Sov. Math. Dokl. **20**(2), 1979, 417-421.
- [15] O.R. Musin, The problem of the twenty-five spheres, Russian Math. Surveys, **58** (2003), 794-795.
- [16] O.R. Musin, The kissing problem in three dimensions, Discrete Comput. Geom., **35** (2006), 375-384.
- [17] O.R. Musin, The kissing number in four dimensions, to appear in Annals of Math. (see math. MG/0309430).
- [18] A.M. Odlyzko and N.J.A. Sloane, New bounds on the number of unit spheres that can touch a unit sphere in n dimensions, J. of Combinatorial Theory A26(1979), 210-214.
- [19] F. Pfender and G.M. Ziegler, Kissing numbers, sphere packings, and some unexpected proofs, Notices Amer. Math. Soc., **51**(2004), 873-883.
- [20] H. Sachs, No more than nine unit balls can touch a closed hemisphere, Studia Sci. Math. Hungar., **21** (1986), 203-206.
- [21] I.J. Schoenberg, Positive definite functions on spheres, Duke Math. J., **9** (1942), 96-107.
- [22] K. Schütte and B.L. van der Waerden, Das Problem der dreizehn Kugeln, Math. Ann. **125** (1953), 325-334.
- [23] L. Szabó, 21-neighbour packing of equal balls in the 4-dimensional Euclidean space, Geom. Dedicata **38** (1991), 193-197.